

YOU'RE INVITED



PLEASE JOIN US FOR AN INTERACTIVE DISCUSSION:

STEVEN BENEDICT, MD

Director
Department of Neurophysiology
Advanced Neurological Associates
Bellevue, Ohio

Wednesday, October 5, 2016

Registration: 5:30 PM ET

**Dinner and
Clinical Discussion:** 6:00 PM ET

Location:
Mitchell's Steakhouse
1408 Polaris Pkwy
Columbus, OH 43240

RSVP: SHAWN MAY
(614) 402-5426
SMAY@LUNDBECK.COM

To register, please contact your Account Manager or complete and return the attached registration form to Lundbeck Speaker Programs.

This is a promotional program sponsored by Lundbeck and is not approved for CME.

In accordance with PhRMA guidelines, guests are not permitted to attend this program and it is intended only for invited, registered health care professionals. Due to limited seating, we suggest you respond as soon as possible.





Registration Form

PLEASE PRINT CLEARLY

*REQUIRED FOR EVENT REGISTRATION

Return this form via fax: **877-337-4408** or email: **support@Lundbeckspeakerprograms.com**.

You may also call our service center if you have any questions: **877-866-7796**.

4483

Date _____ Location _____

Degree* ☐ MD ☐ DO ☐ PA ☐ NP ☐ RN ☐ PharmD ☐ RPH ☐ Case Manager

☐ Other (please specify) _____

First Name* _____ MI _____ Last Name* _____

Gender* ☐ Male ☐ Female

Specialty* _____ Title* _____

Company/Affiliation _____

Business Address* _____

City/State/Zip Code* _____

Email (required for confirmation materials) _____

Phone _____ Fax _____



If you have special needs and wish to discuss provisions that may make your learning experience more effective, please check here ☐

☐ Please send me information on educational programs, products, services, or other information from Lundbeck.



Frictionless is the coordinator for the
ONFI Lundbeck Speaker Programs.

© 2014 Lundbeck. All rights reserved.
ONFI is a registered trademark of Lundbeck CLB-B-00147a 12/2014

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ONFI safely and effectively. See full prescribing information for ONFI.

ONFI® (clobazam) tablets, for oral use, (iv)
ONFI® (clobazam) oral suspension, (iv)

Initial U.S. Approval: 2011

RECENT MAJOR CHANGES

Dosage and Administration:	
Important Administration Instructions (2.3)	3/2013
Warnings and Precautions:	
Serious Dermatological Reactions (5.4)	11/2013

INDICATIONS AND USAGE

ONFI is a benzodiazepine indicated for adjunctive treatment of seizures associated with Lennox-Gastaut syndrome (LGS) in patients 2 years of age or older (1)

DOSAGE AND ADMINISTRATION

- For doses above 5 mg/day administer in two divided doses (2.1)
- Patients ≤30 kg body weight: Initiate at 5 mg daily and titrate as tolerated up to 20 mg daily (2.1)
- Patients >30 kg body weight: Initiate at 10 mg daily and titrate as tolerated up to 40 mg daily (2.1)
- Dosage adjustment needed in following groups:
 - Geriatric patients (2.4, 8.5)
 - Known CYP2C19 poor metabolizers (2.5)
 - Mild or moderate hepatic impairment; no information for severe hepatic impairment (2.7, 8.8)
- Reduce dose, or discontinue drug gradually (2.2)
- Tablets: Administer whole, broken in half along the score, or crush and mix in applesauce. (2.3)
- Measure prescribed amount of oral suspension using provided adapter and dosing syringe (2.3)
- Tablets and Oral suspension: Can be taken with or without food. (2.3)

DOSAGE FORMS AND STRENGTHS

- Tablet: 10 mg and 20 mg with a functional score (3)
- Oral Suspension: 2.5 mg/mL in 120 mL bottles (3)

CONTRAINDICATIONS

None (4)

WARNINGS AND PRECAUTIONS

- Somnolence or Sedation:* Monitor for central nervous system (CNS) depression. Risk may be increased with concomitant use of other CNS depressants. (5.1, 5.2)
- Withdrawal:* Symptoms may occur with rapid dose reduction or discontinuation. Discontinue ONFI gradually. (5.3)
- Physical and Psychological Dependence:* Monitor patients with a history of substance abuse for signs of habituation and dependence (5.5, 9)
- Serious Dermatological Reactions* (including Stevens-Johnson syndrome and toxic epidermal necrolysis): Discontinue ONFI at first sign of rash unless the rash is clearly not drug-related. (5.4)
- Suicidal Behavior and Ideation:* Monitor for suicidal thoughts or behaviors (5.6)

ADVERSE REACTIONS

Adverse reactions that occurred at least 10% more frequently than placebo in any ONFI dose included constipation, somnolence or sedation, pyrexia, lethargy, and drooling (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Lundbeck at 1-800-455-1141 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Drugs metabolized by CYP2D6:* Lower doses of these drugs may be required when used concomitantly with ONFI (7.1)
- Strong or Moderate CYP2C19 Inhibitors:* Dosage adjustment of ONFI may be necessary (7.2)
- Alcohol:* Increases blood levels of clobazam by about 50% (7.3)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 11/2013

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Dosing Information
- 2.2 Gradual Withdrawal
- 2.3 Important Administration Instructions
- 2.4 Dosage Adjustments in Geriatric Patients
- 2.5 Dosage Adjustments in CYP2C19 Poor Metabolizers
- 2.6 Patients with Renal Impairment
- 2.7 Dosage Adjustments in Patients with Hepatic Impairment

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Somnolence or Sedation
- 5.2 Potentiation of Sedation from Concomitant Use with Central Nervous System Depressants
- 5.3 Withdrawal Symptoms
- 5.4 Serious Dermatological Reactions
- 5.5 Physical and Psychological Dependence
- 5.6 Suicidal Behavior and Ideation

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Post Marketing Experience

7 DRUG INTERACTIONS

- 7.1 Effect of ONFI on Other Drugs
- 7.2 Effect of Other Drugs on ONFI
- 7.3 CNS Depressants and Alcohol

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 CYP2C19 Poor Metabolizers
- 8.7 Renal Impairment
- 8.8 Hepatic Impairment

9 DRUG ABUSE AND DEPENDENCE

- 9.1 Controlled Substance
- 9.2 Abuse
- 9.3 Dependence

10 OVERDOSAGE

- 10.1 Signs and Symptoms of Overdosage
- 10.2 Management of Overdosage

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics
- 12.5 Pharmacogenomics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

ONFI® (clobazam) is indicated for the adjunctive treatment of seizures associated with Lennox-Gastaut syndrome (LGS) in patients 2 years of age or older.

2 DOSAGE AND ADMINISTRATION

2.1 Dosing Information

A daily dose of ONFI greater than 5 mg should be administered in divided doses twice daily; a 5 mg daily dose can be administered as a single dose. Dose patients according to body weight. Individualize dosing within each body weight group, based on clinical efficacy and tolerability. Each dose in Table 1 (e.g. 5 to 20 mg in ≤30 kg weight group) has been shown to be effective, although effectiveness increases with increasing dose [see Clinical Studies (14)]. Do not proceed with dose escalation more rapidly than weekly, because serum concentrations of clobazam and its active metabolite require 5 and 9 days, respectively, to reach steady-state.

Table 1. Recommended Total Daily Dosing by Weight Group

	≤30 kg Body Weight	>30 kg Body Weight
Starting Dose	5 mg	10 mg
Starting Day 7	10 mg	20 mg
Starting Day 14	20 mg	40 mg

2.2 Gradual Withdrawal

As with all antiepileptic drugs and benzodiazepines, withdraw ONFI gradually. Taper by decreasing the total daily dose by 5-10 mg/day on a weekly basis until discontinued [see Warnings and Precautions (5.3)].

2.3 Important Administration Instructions

Instruct patients to read the “Instructions for Use” carefully for complete directions on how to properly dose and administer ONFI oral suspension.

ONFI Tablet Oral Administration

ONFI tablets can be taken with or without food.

ONFI tablets can be administered whole, broken in half along the score, or crushed and mixed in applesauce.

ONFI Oral Suspension Oral Administration

ONFI oral suspension can be taken with or without food [see Clinical Pharmacology (12.3)].

Shake ONFI Oral Suspension well before every administration. When administering the oral suspension, use only the oral dosing syringe provided with the product. Each carton includes two syringes, but only one syringe should be used for dosing. The second oral syringe is reserved as a replacement in case the first syringe is damaged or lost. Insert the provided adapter firmly into the neck of the bottle before first use and keep the adapter in place for the duration of the usage of the bottle. To withdraw the dose, insert the dosing syringe into the adapter and invert the bottle then slowly pull back the plunger to prescribed dose. After removing the syringe from the bottle adapter, slowly squirt ONFI Oral Suspension into the corner of the patient’s mouth. Replace the cap after each use. The cap fits over the adapter when the adapter is properly placed. See ONFI Oral Suspension “Instructions for Use” for complete instruction on how to properly dose and administer the ONFI Oral Suspension.

2.4 Dosage Adjustments in Geriatric Patients

Plasma concentrations at any given dose are generally higher in the elderly: proceed slowly with dose escalation. The starting dose should be 5 mg/day for all elderly patients. Then titrate elderly patients according to weight, but to half the dose presented in Table 1, as tolerated. If necessary and based upon clinical response, an additional titration to the maximum dose (20 mg/day or 40 mg/day, depending on weight) may be started on day 21 [see Use in Specific Populations (8.5)].

2.5 Dosage Adjustments in CYP2C19 Poor Metabolizers

In CYP2C19 poor metabolizers, levels of N-desmethyloclobazam, clobazam’s active metabolite, will be increased. Therefore, in patients known to be CYP2C19 poor metabolizers, the starting dose should be 5 mg/day and dose titration should proceed slowly according to weight, but to half the dose presented in Table 1, as tolerated. If necessary and based upon clinical response, an additional titration to the maximum dose (20 mg/day or 40 mg/day, depending on the weight group) may be started on day 21 [see Use in Specific Populations (8.6), Clinical Pharmacology (12.5)].

2.6 Patients with Renal Impairment

No dose adjustment is required for patients with mild and moderate renal impairment. There is no experience with ONFI in patients with severe renal impairment or end stage renal disease (ESRD). It is not known if clobazam or its active metabolite, N-desmethyloclobazam, is dialyzable [see Use in Specific Populations (8.7), Clinical Pharmacology (12.3)].

2.7 Dosage Adjustments in Patients with Hepatic Impairment

ONFI is hepatically metabolized; however, there are limited data to characterize the effect of hepatic impairment on the pharmacokinetics of ONFI. For this reason, proceed slowly with dosing escalations. For patients with mild to moderate hepatic impairment (Child-Pugh score 5-9), the starting dose should be 5 mg/day in both weight groups. Then titrate patients according to weight, but to half the dose presented in Table 1, as tolerated. If necessary and based upon clinical response, start an additional titration on day 21 to the maximum dose (20 mg/day or 40 mg/day, depending on the weight group). There is inadequate information about metabolism of ONFI in patients with severe hepatic impairment. Therefore no dosing recommendation in those patients can be given [see Use in Specific Populations (8.8), Clinical Pharmacology (12.3)].

3 DOSAGE FORMS AND STRENGTHS

Tablets: 10 mg and 20 mg with a functional score for oral administration.
Each ONFI tablet is a white to off-white, oval tablet with a functional score on one side and either a “1” and “0” or a “2” and “0” debossed on the other side.
Oral Suspension: 2.5 mg/mL for oral administration. Each bottle contains 120 mL of an off-white suspension.

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

5.1 Somnolence or Sedation

ONFI causes somnolence and sedation. In clinical trials, somnolence or sedation was reported at all effective doses and was dose-related.
In general, somnolence and sedation begin within the first month of treatment and may diminish with continued treatment. Prescribers should monitor patients for somnolence and sedation, particularly with concomitant use of other central nervous system depressants. Prescribers should caution patients against engaging in hazardous activities requiring mental alertness, such as operating dangerous machinery or motor vehicles, until the effect of ONFI is known.

5.2 Potentiation of Sedation from Concomitant Use with Central Nervous System Depressants

Since ONFI has a central nervous system (CNS) depressant effect, patients or their caregivers should be cautioned against simultaneous use with other CNS depressant drugs or alcohol, and cautioned that the effects of other CNS depressant drugs or alcohol may be potentiated.

5.3 Withdrawal Symptoms

Abrupt discontinuation of ONFI should be avoided. ONFI should be tapered by decreasing the dose every week by 5-10 mg/day until discontinuation [see Dosage and Administration (2.2)].
Withdrawal symptoms occurred following abrupt discontinuation of ONFI; the risk of withdrawal symptoms is greater with higher doses.
As with all antiepileptic drugs, ONFI should be withdrawn gradually to minimize the risk of precipitating seizures, seizure exacerbation, or status epilepticus.
Withdrawal symptoms (e.g., convulsions, psychosis, hallucinations, behavioral disorder, tremor, and anxiety) have been reported following abrupt discontinuance of benzodiazepines. The more severe withdrawal symptoms have usually been limited to patients who received excessive doses over an extended period of time, followed by an abrupt discontinuation. Generally milder withdrawal symptoms (e.g., dysphoria, anxiety, and insomnia) have been reported following abrupt discontinuance of benzodiazepines taken continuously at therapeutic doses for several months.

5.4 Serious Dermatological Reactions

Serious skin reactions, including Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN), have been reported with ONFI in both children and adults during the post-marketing period. Patients should be closely monitored for signs or symptoms of SJS/TEN, especially during the first 8 weeks of treatment initiation or when re-introducing therapy. ONFI should be discontinued at the first sign of rash, unless the rash is clearly not drug-related. If signs or symptoms suggest SJS/TEN, use of this drug should not be resumed and alternative therapy should be considered.

5.5 Physical and Psychological Dependence

Patients with a history of substance abuse should be under careful surveillance when receiving ONFI or other psychotropic agents because of the predisposition of such patients to habituation and dependence [see Drug Abuse and Dependence (9)].

5.6 Suicidal Behavior and Ideation

Antiepileptic drugs (AEDs), including ONFI, increase the risk of suicidal thoughts or behavior in patients taking these drugs for any indication. Patients treated with any AED for any indication should be monitored for the emergence or worsening of depression, suicidal thoughts or behavior, and/or any unusual changes in mood or behavior.
Pooled analyses of 199 placebo-controlled clinical trials (mono- and adjunctive therapy) of 11 different AEDs showed that patients randomized to one of the AEDs had approximately twice the risk (adjusted relative risk 1.8, 95% confidence interval [CI]: 1.2, 2.7) of suicidal thinking or behavior compared to patients randomized to placebo. In these trials, which had a median treatment duration of 12 weeks, the estimated incidence rate of suicidal behavior or ideation among 27,863 AED treated patients was 0.43%, compared to 0.24% among 16,029 placebo treated patients, representing an increase of approximately one case of suicidal thinking or behavior for every 530 patients treated. There were four suicides in drug treated patients in the trials and none in placebo treated patients, but the number is too small to allow any conclusion about drug effect on suicide.
The increased risk of suicidal thoughts or behavior with AEDs was observed as early as one week after starting drug treatment with AEDs and persisted for the duration of treatment assessed. Because most trials included in the analysis did not extend beyond 24 weeks, the risk of suicidal thoughts or behavior beyond 24 weeks could not be assessed.
The risk of suicidal thoughts or behavior was generally consistent among drugs in the data analyzed. The finding of increased risk with AEDs of varying mechanisms of action and across a range of indications suggests that the risk applies to all AEDs used for any indication. The risk did not vary substantially by age (5-100 years) in the clinical trials analyzed. Table 2 shows absolute and relative risk by indication for all evaluated AEDs.

Table 2. Risk by Indication for Antiepileptic Drugs in the Pooled Analysis

Indication	Placebo Patients with Events per 1000 Patients	Drug Patients with Events per 1000 Patients	Relative Risk: Incidence of Drug Events in Drug Patients/Incidence in Placebo Patients	Risk Difference: Additional Drug Patients with Events per 1000 Patients
Epilepsy	1.0	3.4	3.5	2.4
Psychiatric	5.7	8.5	1.5	2.9
Other	1.0	1.8	1.9	0.9
Total	2.4	4.3	1.8	1.9

The relative risk for suicidal thoughts or behavior was higher in clinical trials for epilepsy than in clinical trials for psychiatric or other conditions, but the absolute risk differences were similar for the epilepsy and psychiatric indications.
Anyone considering prescribing ONFI or any other AED must balance the risk of suicidal thoughts or behavior with the risk of untreated illness. Epilepsy and many other illnesses for which AEDs are prescribed are themselves associated with morbidity and mortality and an increased risk of suicidal thoughts and behavior. Should suicidal thoughts and behavior emerge during treatment, the prescriber needs to consider whether the emergence of these symptoms in any given patient may be related to the illness being treated.
Patients, their caregivers, and families should be informed that AEDs increase the risk of suicidal thoughts and behavior and should be advised of the need to be alert for the emergence or worsening of the signs and symptoms of depression, any unusual changes in mood or behavior, or the emergence of suicidal thoughts, behavior, or thoughts about self-harm. Behaviors of concern should be reported immediately to healthcare providers.

6 ADVERSE REACTIONS

Clinically significant adverse reactions that appear in other sections of the labeling include the following:

- Somnolence or Sedation [see Warnings and Precautions (5.1)]
- Potentiation of Sedation from Concomitant Use with Central Nervous System Depressants [see Warnings and Precautions (5.2)]
- Withdrawal Symptoms [see Warnings and Precautions (5.3)]
- Serious Dermatological Reactions [see Warnings and Precautions (5.4)]
- Physical and Psychological Dependence [see Warnings and Precautions (5.5)]
- Suicidal Behavior and Ideation [see Warnings and Precautions (5.6)]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.
During its development for the adjunctive treatment of seizures associated with LGS, ONFI was administered to 333 healthy volunteers and 300 patients with a current or prior diagnosis of LGS, including 197 patients treated for 12 months or more. The conditions and duration of exposure varied greatly and included single- and multiple-dose clinical pharmacology studies in healthy volunteers and two double-blind studies in patients with LGS (Study 1 and 2) [see Clinical Studies (14)]. Only Study 1 included a placebo group, allowing comparison of adverse reaction rates on ONFI at several doses to placebo.
Adverse Reactions Leading to Discontinuation in an LGS Placebo Controlled Clinical Trial (Study 1)
The adverse reactions associated with ONFI treatment discontinuation in ≥1% of patients in decreasing order of frequency included lethargy, somnolence, ataxia, aggression, fatigue, and insomnia.

Most Common Adverse Reactions in an LGS Placebo Controlled Clinical Trial (Study 1)

Table 3 lists the adverse reactions that occurred in ≥5% of ONFI treated patients (at any dose), and at a rate greater than placebo treated patients, in the randomized, double-blind, placebo-controlled, parallel group clinical study of adjunctive AED therapy for 15 weeks (Study 1).

Table 3. Adverse Reactions Reported for ≥5% of Patients and More Frequently than Placebo in Any Treatment Group

	Placebo N=59 %	ONFI Dose Level			All ONFI N=179 %
		Low ^a N=58 %	Medium ^b N=62 %	High ^c N=59 %	
Gastrointestinal Disorders					
Vomiting	5	9	5	7	7
Constipation	0	2	2	10	5
Dysphagia	0	0	0	5	2
General Disorders and Administration Site Conditions					
Pyrexia	3	17	10	12	13
Irritability	5	3	11	5	7
Fatigue	2	5	5	3	5
Infections and Infestations					
Upper respiratory tract infection	10	10	13	14	12
Pneumonia	2	3	3	7	4
Urinary tract infection	0	2	5	5	4
Bronchitis	0	2	0	5	2
Metabolism and Nutrition Disorders					
Decreased appetite	3	3	0	7	3
Increased appetite	0	2	3	5	3
Nervous System Disorders					
Somnolence or Sedation	15	17	27	32	26
Somnolence	12	16	24	25	22
Sedation	3	2	3	9	5
Lethargy	5	10	5	15	10
Drooling	3	0	13	14	9
Ataxia	3	3	2	10	5
Psychomotor hyperactivity	3	3	3	5	4
Dysarthria	0	2	2	5	3
Psychiatric Disorders					
Aggression	5	3	8	14	8
Insomnia	2	2	5	7	5
Respiratory Disorders					
Cough	0	3	5	7	5

^aMaximum daily dose of 5 mg for ≤30 kg body weight; 10 mg for >30 kg body weight
^bMaximum daily dose of 10 mg for ≤30 kg body weight; 20 mg for >30 kg body weight
^cMaximum daily dose of 20 mg for ≤30 kg body weight; 40 mg for >30 kg body weight

6.2 Post Marketing Experience

These reactions are reported voluntarily from a population of uncertain size; therefore, it is not possible to estimate their frequency or establish a causal relationship to drug exposure. Adverse reactions are categorized by system organ class.

Blood Disorders: Anemia, eosinophilia, leukopenia, thrombocytopenia

Eye Disorders: Diplopia, vision blurred

Gastrointestinal Disorders: Abdominal distention

Investigations: Hepatic enzyme increased

Musculoskeletal: Muscle spasms

Psychiatric Disorders: Agitation, anxiety, apathy, confusional state, depression, delirium, delusion, hallucination

Respiratory Disorders: Aspiration, respiratory depression

Skin and Subcutaneous Tissue Disorders: Rash, urticaria

7 DRUG INTERACTIONS

7.1 Effect of ONFI on Other Drugs

Hormonal Contraceptives

ONFI is a weak CYP3A4 inducer. As some hormonal contraceptives are metabolized by CYP3A4, their effectiveness may be diminished when given with ONFI. Additional non-hormonal forms of contraception are recommended when using ONFI [see Clinical Pharmacology (12.3), Patient Counseling Information (17)].

Drugs Metabolized by CYP2D6

ONFI inhibits CYP2D6. Dose adjustment of drugs metabolized by CYP2D6 may be necessary [see Clinical Pharmacology (12.3)].

7.2 Effect of Other Drugs on ONFI

Strong and moderate inhibitors of CYP2C19

Strong and moderate inhibitors of CYP2C19 may result in increased exposure to N-desmethyloclobazam, the active metabolite of clobazam. This may increase the risk of dose-related adverse reactions. Dosage adjustment of ONFI may be necessary when co-administered with strong CYP2C19 inhibitors (e.g., fluconazole, fluvoxamine, ticlopidine) or moderate CYP2C19 inhibitors (e.g., omeprazole) [see Clinical Pharmacology (12.3)].

7.3 CNS Depressants and Alcohol

Concomitant use of ONFI with other CNS depressants may increase the risk of sedation and somnolence [see Warnings and Precautions (5.2)].

Alcohol, as a CNS depressant, will interact with ONFI in a similar way and also increases clobazam’s maximum plasma exposure by approximately 50%. Therefore, caution patients or their caregivers against simultaneous use with other CNS depressant drugs or alcohol, and caution that the effects of other CNS depressant drugs or alcohol may be potentiated [see Warnings and Precautions (5.2)].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Registry: To provide information regarding the effects of *in utero* exposure to ONFI, physicians are advised to recommend that pregnant patients taking ONFI enroll in the North American Antiepileptic Drug (NAAED) Pregnancy Registry. This can be done by calling the toll free number 1-888-233-2334, and must be done by patients themselves or their caregiver. Information on the registry can also be found at the website <http://www.aedpregnancyregistry.org/>.

Pregnancy Category C.

There are no adequate and well-controlled studies of ONFI in pregnant women and no adequate developmental toxicity studies of clobazam in animals.

Although limited, the available animal data suggest developmental toxicity, including an increased incidence of fetal abnormalities following oral administration of clobazam to pregnant animals at doses similar to those used clinically.

Data for other benzodiazepines suggest the possibility of adverse effects in animals and humans. Long-term effects on neurobehavioral and immunological function have been reported in rodents following prenatal exposure to benzodiazepines. Neonatal flaccidity, respiratory and feeding difficulties, hypothermia, and withdrawal symptoms have been reported in infants born to mothers who received benzodiazepines, including clobazam, late in pregnancy.

Therefore, ONFI should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

8.3 Nursing Mothers

ONFI is excreted in human milk. The effects of this exposure on infants are unknown.

8.4 Pediatric Use

The safety and effectiveness in patients less than 2 years of age have not been established.

In a study in which clobazam (4, 36, or 120 mg/kg/day) was orally administered to rats during the juvenile period of development (postnatal days 14 to 48), adverse effects on growth (decreased bone density and bone length) and behavior (altered motor activity and auditory startle response; learning deficit) were observed at the high dose. The effect on bone density, but not on behavior, was reversible when drug was discontinued. The no-effect level for juvenile toxicity (36 mg/kg/day) was associated with plasma exposures (AUC) to clobazam and its major active metabolite, N-desmethyloclobazam, less than those expected at therapeutic doses in pediatric patients.

8.5 Geriatric Use

Clinical studies of ONFI did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. However, elderly subjects appear to eliminate clobazam more slowly than younger subjects based on population pharmacokinetic analysis. For these reasons, the initial dose in elderly patients should be 5 mg/day. Patients should be titrated initially to 10-20 mg/day. Patients may be titrated further to a maximum daily dose of 40 mg if tolerated [see Dosage and Administration (2.4), Clinical Pharmacology (12.3)].

8.6 CYP2C19 Poor Metabolizers

Concentrations of clobazam’s active metabolite, N-desmethyloclobazam, are higher in CYP2C19 poor metabolizers than in extensive metabolizers. For this reason, dosage modification is recommended [see Dosage and Administration (2.5), Clinical Pharmacology (12.3)].

8.7 Renal Impairment

The pharmacokinetics of ONFI were evaluated in patients with mild and moderate renal impairment. There were no significant differences in systemic exposure (AUC and C_{max}) between patients with mild or moderate renal impairment and healthy subjects. No dose adjustment is required for patients with mild and moderate renal impairment. There is essentially no experience with ONFI in patients with severe renal impairment or ESRD. It is not known if clobazam or its active metabolite, N-desmethyloclobazam, is dialyzable [see Dosage and Administration (2.6), Clinical Pharmacology (12.3)].

8.8 Hepatic Impairment

ONFI is hepatically metabolized; however, there are limited data to characterize the effect of hepatic impairment on the pharmacokinetics of ONFI. For this reason, dosage adjustment is recommended in patients with mild to moderate hepatic impairment (Child-Pugh score 5-9). There is inadequate information about metabolism of ONFI in patients with severe hepatic impairment [see Dosage and Administration (2.7), Clinical Pharmacology (12.3)].

9 DRUG ABUSE AND DEPENDENCE

9.1 Controlled Substance

ONFI contains clobazam which is a Schedule IV controlled substance.

9.2 Abuse

ONFI can be abused in a similar manner as other benzodiazepines, such as diazepam.

The pharmacological profile of ONFI is similar to that of other benzodiazepines listed in Schedule IV of the Controlled Substance Act, particularly in its potentiation of GABAergic transmission through its action on GABA_A receptors, which leads to sedation and somnolence.

The World Health Organization epidemiology database contains reports of drug abuse, misuse, and overdoses associated with clobazam.

Drug abuse is the intentional non-therapeutic use of a drug, repeatedly or even sporadically, for its rewarding psychological or physiological effects.

9.3 Dependence

Dependence

Physical dependence is a state of adaptation that is manifested by a specific withdrawal syndrome that can be produced by abrupt cessation, rapid dose reduction, decreasing blood levels of the drug, and/or administration of an antagonist. In clinical trials, cases of dependency were reported following abrupt discontinuation of ONFI.

The risk of dependence is present even with use of ONFI at the recommended dose range over periods of only a few weeks. The risk of dependence increases with increasing dose and duration of treatment. The risk of dependence is increased in patients with a history of alcohol or drug abuse.

Withdrawal

Abrupt discontinuation of ONFI causes withdrawal symptoms. As with other benzodiazepines, ONFI should be withdrawn gradually [see Dosage and Administration (2.2), Warnings and Precautions (5.3)].

In ONFI clinical pharmacology trials in healthy volunteers, the most common withdrawal symptoms after abrupt discontinuation were headache, tremor, insomnia, anxiety, irritability, drug withdrawal syndrome, palpitations, and diarrhea [see Warnings and Precautions (5.3)].

Other withdrawal reactions to clobazam reported in the literature include restlessness, panic attacks, profuse sweating, difficulty in concentrating, nausea and dry retching, weight loss, blurred vision, photophobia, and muscle pain and stiffness. In general, benzodiazepine withdrawal may cause seizures, psychosis, and hallucinations [see Warnings and Precautions (5.3)].

10 OVERDOSAGE

10.1 Signs and Symptoms of Overdosage

Overdose and intoxication with benzodiazepines, including ONFI, may lead to CNS depression, associated with drowsiness, confusion and lethargy, possibly progressing to ataxia, respiratory depression, hypotension, and, rarely, coma or death. The risk of a fatal outcome is increased in cases of combined poisoning with other CNS depressants, including alcohol.


10.2 Management of Overdosage

The management of ONFI overdose may include gastric lavage and/or administration of activated charcoal, intravenous fluid replenishment, early control of airway and general supportive measures, in addition to monitoring level of consciousness and vital signs. Hypotension can be treated by replenishment with plasma substitutes and, if necessary, with sympathomimetic agents.

The efficacy of supplementary administration of physostigmine (a cholinergic agent) or of flumazenil (a benzodiazepine antagonist) in ONFI overdose has not been assessed. The administration of flumazenil in cases of benzodiazepine overdose can lead to withdrawal and adverse reactions. Its use in patients with epilepsy is typically not recommended.

11 DESCRIPTION

Table 4. Description

Proprietary Name:	ONFI®
Established Name:	Clobazam
Dosage Forms:	Tablet and Oral Suspension
Route of Administration:	Oral
Established Pharmacologic Class of Drug:	Benzodiazepine
Chemical Name:	7-Chloro-1-methyl-5-phenyl-1H-1,5 benzodiazepine-2,4(3H,5H)-dione
Structural Formula:	

Clobazam is a white or almost white, crystalline powder with a slightly bitter taste; is slightly soluble in water, sparingly soluble in ethanol, and freely soluble in methylene chloride. The melting range of clobazam is from 182°C to 185°C. The molecular formula is C₁₆H₁₃O₂N₂Cl and the molecular weight is 300.7.

Each ONFI tablet contains 10 mg or 20 mg of clobazam. Tablets also contain as inactive ingredients: corn starch, lactose monohydrate, magnesium stearate, silicon dioxide, and talc.

ONFI is also available for oral administration as an off-white suspension containing clobazam at a concentration of 2.5 mg/mL. Inactive ingredients include magnesium aluminum silicate, xanthan gum, citric acid monohydrate, disodium hydrogen phosphate dihydrate, simethicone emulsion, polysorbate 80, methylparaben, propylparaben, propylene glycol, sucralose, maltitol solution, berry flavor, purified water.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The exact mechanism of action for clobazam, a 1,5-benzodiazepine, is not fully understood but is thought to involve potentiation of GABAergic neurotransmission resulting from binding at the benzodiazepine site of the GABA_A receptor.

12.2 Pharmacodynamics

Effects on Electrocardiogram

The effect of ONFI 20 mg and 80 mg administered twice daily on QTc interval was evaluated in a randomized, evaluator blinded, placebo-, and active-controlled (moxifloxacin 400 mg) parallel thorough QT study in 280 healthy subjects. In a study with demonstrated ability to detect small effects, the upper bound of the one-sided 95% confidence interval for the largest placebo adjusted, baseline-corrected QTc based on the Fridericia correction method was below 10 ms, the threshold for regulatory concern. Thus, at a dose two times the maximum recommended dose, ONFI did not prolong the QTc interval to any clinically relevant extent.

12.3 Pharmacokinetics

The peak plasma levels (C_{max}) and the area under the curve (AUC) of clobazam are dose-proportional over the dose range of 10-80 mg following single- or multiple-dose administration of ONFI. Based on a population pharmacokinetic analysis, the pharmacokinetics of clobazam are linear from 5-160 mg/day. Clobazam is converted to N-desmethyclobazam which has about 1/5 the activity of clobazam. The estimated mean elimination half-lives (t_{1/2}) of clobazam and N-desmethyclobazam were 36-42 hours and 71-82 hours, respectively.

Absorption

Clobazam is rapidly and extensively absorbed following oral administration. The time to peak concentrations (T_{max}) of clobazam tablets under fasted conditions ranged from 0.5 to 4 hours after single- or multiple-dose administrations. The relative bioavailability of clobazam tablets compared to an oral solution is approximately 100%. After single dose administration of the oral suspension under fasted conditions, the T_{max} ranged from 0.5 to 2 hours. Based on exposure (C_{max} and AUC) of clobazam, ONFI tablets and suspension were shown to have similar bioavailability under fasted conditions. The administration of ONFI tablets with food or when crushed in applesauce does not affect absorption. Although not studied, the oral bioavailability of the oral suspension is unlikely to be affected under fed conditions.

Distribution

Clobazam is lipophilic and distributes rapidly throughout the body. The apparent volume of distribution at steady state was approximately 100 L. The *in vitro* plasma protein binding of clobazam and N-desmethyclobazam is approximately 80-90% and 70%, respectively.

Metabolism and Excretion

Clobazam is extensively metabolized in the liver, with approximately 2% of the dose recovered in urine and 1% in feces as unchanged drug. The major metabolic pathway of clobazam involves N-demethylation, primarily by CYP3A4 and to a lesser extent by CYP2C19 and CYP2B6. N-desmethyclobazam, an active metabolite, is the major circulating metabolite in humans, and at therapeutic doses, plasma concentrations are 3-5 times higher than those of the parent compound. Based on animal and *in vitro* receptor binding data, estimates of the relative potency of N-desmethyclobazam compared to parent compound range from 1/5 to equal potency. N-desmethyclobazam is extensively metabolized, mainly by CYP2C19. N-desmethyclobazam and its metabolites comprise ~94% of the total drug-related components in urine. Following a single oral dose of radiolabeled drug, approximately 11% of the dose was excreted in the feces and approximately 82% was excreted in the urine.

The polymorphic CYP2C19 is the major contributor to the metabolism of the pharmacologically active N-desmethyclobazam [see Clinical Pharmacology (12.5)]. In CYP2C19 poor metabolizers, levels of N-desmethyclobazam were 5-fold higher in plasma and 2- to 3-fold higher in the urine than in CYP2C19 extensive metabolizers.

Pharmacokinetics in Specific Populations

Age

Population pharmacokinetic analyses showed that the clearance of clobazam is lower in elderly subjects compared to other age groups (ages 2 to 64). Dosing should be adjusted in the elderly [see Dosage and Administration (2.4)].

Sex

Population pharmacokinetic analyses showed no difference in the clearance of clobazam between women and men.

Race

Population pharmacokinetic analyses including Caucasian (75%), African American (15%), and Asian (9%) subjects showed that there is no evidence of clinically significant effect of race on the clearance of clobazam.

Renal Impairment

The effect of renal impairment on the pharmacokinetics of clobazam was evaluated in patients with mild (creatinine clearance [CL_{CR}] >50 to 80 mL/min; N=6) and moderate (CL_{CR}=30 to 50 mL/min; N=6) renal dysfunction, with matching healthy controls (N=6), following administration of multiple

doses of ONFI 20 mg/day. There were insignificant changes in C_{max} (3-24%) and AUC (≤13%) for clobazam or N-desmethyclobazam in patients with mild or moderate renal impairment compared to patients with normal renal function. Patients with severe renal impairment or ESRD were not included in this study.

Hepatic Impairment

There are limited data to characterize the effect of hepatic impairment on the pharmacokinetics of clobazam. In a small study, the pharmacokinetics of a 20 mg single oral dose of ONFI in 9 patients with liver impairment were compared to healthy controls (N=6). The C_{max} and the mean plasma clearance of clobazam, as well as the C_{max} of N-desmethyclobazam, showed no significant change compared to the healthy controls. The AUC values of N-desmethyclobazam in these patients were not available. Adjust dosage in patients with hepatic impairment [see Dosage and Administration (2.7)].

Drug Interaction Studies

In vitro studies:

Clobazam did not inhibit CYP1A2, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP3A4, UGT1A1, UGT1A4, UGT1A6, or UGT2B4 *in vitro*. N-desmethyclobazam showed weak inhibition of CYP2C9, UGT1A4, UGT1A6 and UGT2B4.

Clobazam and N-desmethyclobazam did not significantly increase CYP1A2 or CYP2C19 activities, but did induce CYP3A4 activity in a concentration-dependent manner. Clobazam and N-desmethyclobazam also increased UGT1A1 mRNA but at concentrations much higher than therapeutic levels. The potential for clobazam or N-desmethyclobazam to induce CYP2B6 and CYP2C8 has not been evaluated.

Clobazam and N-desmethyclobazam do not inhibit P-glycoprotein (P-gp), but are P-gp substrates.

In vivo studies:

Potential for ONFI to Affect Other Drugs

The effect of repeated 40 mg once-daily doses of ONFI on the pharmacokinetic profiles of single-dose dextromethorphan (CYP2D6 substrate), midazolam (CYP3A4 substrate), caffeine (CYP1A2 substrate), and tolbutamide (CYP2C9 substrate), was studied when these probe substrates were given as a drug cocktail (N=18).

Clobazam increased AUC and C_{max} of dextromethorphan by 90% and 59%, respectively, reflecting its inhibition of CYP2D6 *in vivo*. Drugs metabolized by CYP2D6 may require dose adjustment when used with ONFI.

Clobazam decreased the AUC and C_{max} of midazolam by 27% and 24%, respectively, and increased the AUC and C_{max} of the metabolite 1-hydroxymidazolam by 4-fold and 2-fold, respectively. This level of induction does not call for dosage adjustment of drugs that are primarily metabolized by CYP3A4 when used concomitantly with ONFI. Some hormonal contraceptives are metabolized by CYP3A4 and their effectiveness may be diminished when given with ONFI [see Drug Interactions (7.1)]. Repeated ONFI doses had no effect on caffeine and tolbutamide.

A population pharmacokinetic analysis indicated clobazam did not affect the exposure of valproic acid (a CYP2C9/2C19 substrate) or lamotrigine (a UGT substrate).

Potential for Other Drugs to Affect ONFI

Co-administration of ketoconazole (a strong CYP3A4 inhibitor) 400 mg once-daily for 5 days increased clobazam AUC by 54%, with an insignificant effect on clobazam C_{max}. There was no significant change in AUC and C_{max} of N-desmethyclobazam (N=18).

Strong (e.g., fluconazole, fluvoxamine, ticlopidine) and moderate (e.g., omeprazole) inhibitors of CYP2C19 may result in up to a 5-fold increase in exposure to N-desmethyclobazam, the active metabolite of clobazam, based on extrapolation from pharmacogenomic data [see Clinical Pharmacology (12.5)]. Dosage adjustment of ONFI may be necessary when co-administered with strong or moderate CYP2C19 inhibitors [see Drug Interactions (7.2)].

The effects of concomitant antiepileptic drugs that are CYP3A4 inducers (phenobarbital, phenytoin, and carbamazepine), CYP2C9 inducers (valproic acid, phenobarbital, phenytoin, and carbamazepine), and CYP2C9 inhibitors (felbamate and oxcarbazepine) were evaluated using data from clinical trials. Results of population pharmacokinetic analysis show that these concomitant antiepileptic drugs did not significantly alter the pharmacokinetics of clobazam or N-desmethyclobazam at steady-state.

Alcohol has been reported to increase the maximum plasma exposure of clobazam by approximately 50%. Alcohol may have additive CNS depressant effects when taken with ONFI [see Warnings and Precautions (5.2), Drug Interactions (7.3)].

12.5 Pharmacogenomics

The polymorphic CYP2C19 is the main enzyme that metabolizes the pharmacologically active N-desmethyclobazam. Compared to CYP2C19 extensive metabolizers, N-desmethyclobazam AUC and C_{max} are approximately 3-5 times higher in poor metabolizers (e.g., subjects with *2/*2 genotype) and 2 times higher in intermediate metabolizers (e.g., subjects with *1/*2 genotype). The prevalence of CYP2C19 poor metabolism differs depending on racial/ethnic background. Dosage in patients who are known CYP2C19 poor metabolizers may need to be adjusted [see Dosage and Administration (2.5)].

The systemic exposure of clobazam is similar for both CYP2C19 poor and extensive metabolizers.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility Carcinogenesis

The carcinogenic potential of clobazam has not been adequately assessed.

In a limited study in rats, oral administration of clobazam (4, 20, and 100 mg/kg/day) for 2 years resulted in an increased incidence of thyroid follicular cell adenomas in males at the high dose.

Mutagenesis

Clobazam and the major active metabolite, N-desmethyclobazam, were negative for genotoxicity, based on data from a battery of *in vitro* (bacteria reverse mutation, mammalian clastogenicity) and *in vivo* (mouse micronucleus) assays.

Impairment of Fertility

There are no adequate studies of the effects of clobazam on fertility.

14 CLINICAL STUDIES

The effectiveness of ONFI for the adjunctive treatment of seizures associated with Lennox-Gastaut syndrome was established in two multicenter controlled studies (Study 1 and Study 2). Both studies were similar in terms of disease characteristics and concomitant AED treatments. The most common concomitant AED treatments at baseline included: valproate, lamotrigine, levetiracetam, and topiramate.

Study 1

Study 1 (N=238) was a randomized, double-blind, placebo-controlled study consisting of a 4-week baseline period followed by a 3-week titration period and 12-week maintenance period. Patients age 2-54 years with a current or prior diagnosis of LGS were stratified into 2 weight groups (12.5 kg to ≤30 kg or >30 kg) and then randomized to placebo or one of three target maintenance doses of ONFI according to Table 5.

Table 5. Study 1 Total Daily Dose

	≤30 kg Body Weight	>30 kg Body Weight
Low Dose	5 mg daily	10 mg daily
Medium Dose	10 mg daily	20 mg daily
High Dose	20 mg daily	40 mg daily

Doses above 5 mg/day were administered in two divided doses.

The primary efficacy measure was the percent reduction in the weekly frequency of drop seizures (atonic, tonic, or myoclonic), also known as drop attacks, from the 4-week baseline period to 12-week maintenance period.

The pre-dosing baseline mean weekly drop seizure frequency was 98, 100, 61, and 105 for the placebo, low-, medium-, and high-dose groups, respectively. Figure 1 presents the mean percent reduction in weekly drop seizures from this baseline. All dose groups of ONFI were statistically superior (p≤0.05) to the placebo group. This effect appeared to be dose dependent.

Figure 1. Mean Percent Reduction from Baseline in Weekly Drop Seizure Frequency (Study 1)





Figure 2 shows changes from baseline in weekly drop seizure frequency by category for patients treated with ONFI and placebo in Study 1. Patients in whom the seizure frequency increased are shown at left as “worse.” Patients in whom the seizure frequency decreased are shown in five categories.

Figure 2. Drop Seizure Response by Category for ONFI and Placebo (Study 1)



There was no evidence that tolerance to the therapeutic effect of ONFI developed during the 3-month maintenance period.

Study 2

Study 2 (N=68) was a randomized, double-blind comparison study of high- and low-dose ONFI, consisting of a 4-week baseline period followed by a 3-week titration period and 4-week maintenance period. Patients age 2-25 years with a current or prior diagnosis of LGS were stratified by weight, then randomized to either a low or high dose of ONFI, and then entered a 3-week titration period.

The primary efficacy measure was the percent reduction in the weekly frequency of drop seizures (atonic, tonic, or myoclonic), also known as drop attacks, from the 4-week baseline period to the 4-week maintenance period.

A statistically significantly greater reduction in seizure frequency was observed in the high-dose group compared to the low-dose group (median percent reduction of 93% vs 29%; $p < 0.05$).

16 HOW SUPPLIED/STORAGE AND HANDLING

Each ONFI tablet contains 10 mg or 20 mg of clobazam and is a white to off-white, oval tablet with a functional score on one side and either a “1” and “0” or a “2” and “0” debossed on the other side.

NDC 67386-314-01: 10 mg scored tablet, Bottles of 100

NDC 67386-315-01: 20 mg scored tablet, Bottles of 100

ONFI oral suspension is a berry flavored off-white liquid supplied in a bottle with child-resistant closure. The oral suspension is packaged with a dispenser set which contains two calibrated oral dosing syringes and bottle adapter. Store the oral suspension in an upright position. Use within 90 days of first opening the bottle, then discard any remainder.

NDC 67386-313-21: Bottle containing 120 mL of suspension

Store tablets and oral suspension at 20°C to 25°C (68°F to 77°F). See USP controlled room temperature.

17 PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling (Medication Guide and Instructions for Use).

Inform patients or caregivers of the availability of a Medication Guide and instruct them to read the Medication Guide prior to initiating treatment with ONFI and with each prescription refill. Review the ONFI Medication Guide with every patient or caregiver prior to initiation of treatment. Instruct patients or caregivers that ONFI should be taken only as prescribed.

Somnolence or Sedation

Advise patients or caregivers to check with their healthcare provider before ONFI is taken with other CNS depressants such as other benzodiazepines, opioids, tricyclic antidepressants, sedating antihistamines, or alcohol [see *Warnings and Precautions* (5.1, 5.2)].

If applicable, caution patients about operating hazardous machinery, including automobiles, until they are reasonably certain that ONFI does not affect them adversely (e.g., impair judgment, thinking or motor skills).

Increasing or Decreasing the ONFI Dose

Inform patients or caregivers to consult their healthcare provider before increasing the ONFI dose or abruptly discontinuing ONFI. Advise patients or caregivers that abrupt withdrawal of AEDs may increase their risk of seizure [see *Dosage and Administration* (2.2), *Warnings and Precautions* (5.3)].

Interactions with Hormonal Contraceptives

Counsel women to also use non-hormonal methods of contraception when ONFI is used with hormonal contraceptives and to continue these alternative methods for 28 days after discontinuing ONFI to ensure contraceptive reliability [see *Drug Interactions* (7.1), *Clinical Pharmacology* (12.3)].

Serious Dermatological Reactions

Advise patients or caregivers that serious skin reactions have been reported in patients taking ONFI. Serious skin reactions, including SJS/TEN, may need to be treated in a hospital and may be life-threatening. If a skin reaction occurs while taking ONFI, patients or caregivers should consult with healthcare providers immediately [see *Warnings and Precautions* (5.4)].

Suicidal Thinking and Behavior

Counsel patients, their caregivers, and their families that AEDs, including ONFI, may increase the risk of suicidal thoughts and behavior and advise them of the need to be alert for the emergence or worsening of symptoms of depression, any unusual changes in mood or behavior, or the emergence of suicidal thoughts, behavior, or thoughts of self-harm. Patients should report behaviors of concern immediately to healthcare providers [see *Warnings and Precautions* (5.6)].

Use in Pregnancy

Instruct patients to notify their healthcare provider if they become pregnant or intend to become pregnant during therapy.

Encourage patients to enroll in the NAAED Pregnancy Registry if they become pregnant. This registry is collecting information about the safety of antiepileptic drugs during pregnancy. To enroll, patients can call the toll-free number 1-888-233-2334. Information on the registry can also be found at the website <http://www.aedpregnancyregistry.org> [see *Use in Specific Populations* (8.1)].

Use in Nursing

Instruct patients to notify their physician if they are breast feeding or intend to breast feed during therapy [see *Use in Specific Populations* (8.3)].

Tablets manufactured by: Catalent Pharma Solutions, LLC
Winchester, KY 40391, U.S.A.

Oral suspension manufactured by: Rosemont Pharmaceuticals, Ltd.
Leeds, West Yorkshire LS11 9XE, U.K.

For: Lundbeck
Deerfield, IL 60015, U.S.A.

MEDICATION GUIDE
ONFI® (ON-fee)
(clobazam)
Tablets and Oral Suspension

Read this Medication Guide before you start taking ONFI and each time you get a refill. There may be new information. This information does not take the place of talking to your healthcare provider about your medical condition or treatment.

What is the most important information I should know about ONFI?

Do not stop taking ONFI without first talking to your healthcare provider.

Stopping ONFI suddenly can cause serious problems.

ONFI can cause serious side effects, including:

1. ONFI can make you sleepy or dizzy, slow your thinking, and make you clumsy which may get better over time.

- Do not drive, operate heavy machinery, or do other dangerous activities until you know how ONFI affects you.
- Do not drink alcohol or take other drugs that may make you sleepy or dizzy while taking ONFI until you talk to your healthcare provider. When taken with alcohol or drugs that cause sleepiness or dizziness, ONFI may make your sleepiness or dizziness much worse.

2. ONFI can cause withdrawal symptoms.

- Do not stop taking ONFI all of a sudden without first talking to a healthcare provider. Stopping ONFI suddenly can cause seizures that will not stop (status epilepticus), hearing or seeing things that are not there (hallucinations), shaking, nervousness, and stomach and muscle cramps.
- Talk to your healthcare provider about slowly stopping ONFI to avoid withdrawal symptoms.

3. ONFI can be abused and cause dependence.

- Physical dependence is not the same as drug addiction. Your healthcare provider can tell you more about the differences between physical dependence and drug addiction.

ONFI is a federally controlled substance (C-IV) because it can be abused or lead to dependence. Keep ONFI in a safe place to prevent misuse and abuse. Selling or giving away ONFI may harm others, and is against the law. Tell your healthcare provider if you have ever abused or been dependent on alcohol, prescription medicines or street drugs.

4. Serious skin reactions have been seen with ONFI and may require stopping its use. Do not stop taking ONFI without first talking to your healthcare provider.

- A serious skin reaction can happen at any time during your treatment with ONFI, but is more likely to happen within the first 8 weeks of treatment. These skin reactions need to be treated right away.
- Call your healthcare provider immediately if you have skin blisters, peeling rash, sores in the mouth, hives or any other allergic reaction.

5. Like other antiepileptic drugs, ONFI may cause suicidal thoughts or actions in a very small number of people, about 1 in 500.

Call your healthcare provider right away if you have any of these symptoms, especially if they are new, worse, or worry you:

- thoughts about suicide or dying
- attempts to commit suicide
- new or worse depression
- new or worse anxiety
- feeling agitated or restless
- panic attacks
- trouble sleeping (insomnia)
- new or worse irritability
- acting aggressive, being angry, or violent
- acting on dangerous impulses
- an extreme increase in activity and talking (mania)
- other unusual changes in behavior or mood

How can I watch for early symptoms of suicidal thoughts and actions?

- Pay attention to any changes, especially sudden changes, in mood, behaviors, thoughts, or feelings.

- Keep all follow-up visits with your healthcare provider as scheduled.

Call your healthcare provider between visits as needed, especially if you are worried about symptoms.

Suicidal thoughts or actions can be caused by things other than medicines. If you have suicidal thoughts or actions, your healthcare provider may check for other causes.

What is ONFI?

ONFI is a prescription medicine used along with other medicines to treat seizures associated with Lennox-Gastaut syndrome in people 2 years of age or older.

It is not known if ONFI is safe and effective in children less than 2 years old.

What should I tell my healthcare provider before taking ONFI?

Before you take ONFI, tell your healthcare provider if you:

- have liver or kidney problems
- have lung problems (respiratory disease)
- have or have had depression, mood problems, or suicidal thoughts or behavior
- have any other medical conditions
- use birth control medicine. ONFI may cause your birth control medicine to be less effective. Talk to your healthcare provider about the best birth control method to use.
- are pregnant or plan to become pregnant. **ONFI may harm your unborn baby.**
 - Tell your healthcare provider right away if you become pregnant while taking ONFI. You and your healthcare provider will decide if you should take ONFI while you are pregnant.
 - Children born to mothers receiving benzodiazepine medications (including ONFI) late in pregnancy may be at some risk of experiencing breathing problems, feeding problems, dangerously low body temperature, and withdrawal symptoms.
- If you become pregnant while taking ONFI, talk to your healthcare provider about registering with the North American Antiepileptic Drug Pregnancy Registry. You can register by calling 1-888-233-2334. For more information about the registry go to <http://www.aedpregnancyregistry.org>. The purpose of this registry is to collect information about the safety of antiepileptic drugs during pregnancy.
- ONFI can pass into breast milk. Talk to your healthcare provider about the best way to feed your baby if you take ONFI. You and your healthcare provider should decide if you will take ONFI or breast feed. You should not do both.

Tell your healthcare provider about all the medicines you take, including prescription and nonprescription medicines, vitamins, and herbal supplements. Taking ONFI with certain other medicines can cause side effects or affect how well ONFI or the other medications work. Do not start or stop other medicines without talking to your healthcare provider.

Know the medicines you take. Keep a list of them and show it to your healthcare provider and pharmacist when you get a new medicine.

How should I take ONFI?

- Take ONFI exactly as your healthcare provider tells you to take it.
- Your healthcare provider will tell you how much ONFI to take and when to take it.
- ONFI tablets can be taken whole, broken in half along the score, or crushed and mixed in applesauce.
- ONFI tablets and oral suspension can be taken with or without food.
- **Shake the bottle of ONFI oral suspension well right before you take each dose.**
- Measure your dose of ONFI oral suspension using the bottle adapter and dosing syringes that come with your ONFI oral suspension.
- Read the **Instructions for Use** at the end of this Medication Guide for information on the right way to use ONFI oral suspension.
- Your healthcare provider may change your dose if needed. Do not change your dose of ONFI without talking to your healthcare provider.
- Do not stop taking ONFI without first talking to your healthcare provider.
- Stopping ONFI suddenly can cause serious problems.
- If you take too much ONFI, call your healthcare provider or go to the nearest hospital emergency room right away.

What should I avoid while taking ONFI?

- Do not drive, operate heavy machinery, or do other dangerous activities until you know how ONFI affects you.
- Do not drink alcohol or take other medicines that may make you sleepy or dizzy while taking ONFI until you talk to your healthcare provider. When taken with alcohol or medicines that cause sleepiness or dizziness, ONFI may make your sleepiness or dizziness much worse.

What are the possible side effects of ONFI?

ONFI may cause serious side effects, including:

See “What is the most important information I should know about ONFI?”

The most common side effects of ONFI include:

- | | | |
|----------------|--|---------------------------|
| • sleepiness | • pain with urination | • difficulty sleeping |
| • drooling | • fever | • slurred speech |
| • constipation | • acting aggressive, being angry, or violent | • tiredness |
| • cough | | • problems with breathing |

These are not all the possible side effects of ONFI. For more information, ask your healthcare provider or pharmacist.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store ONFI?

- Store ONFI tablets and oral suspension between 68°F to 77°F (20°C to 25°C).

Tablets

- Keep ONFI tablets in a dry place

Oral Suspension

- Replace the cap securely after opening.
- Keep ONFI oral suspension in an upright position.
- Use ONFI oral suspension within 90 days of first opening the bottle.
- After 90 days safely throw away any ONFI oral suspension that has not been used.

Keep ONFI and all medicines out of the reach of children.

General Information about the safe and effective use of ONFI.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use ONFI for a condition for which it was not prescribed. Do not give ONFI to other people, even if they have the same symptoms that you have. It may harm them.

This Medication Guide summarizes the most important information about ONFI. If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about ONFI that is written for health professionals.

For more information about ONFI, go to www.lundbeckus.com or call Lundbeck at 1-888-514-5204.

What are the ingredients in ONFI?

Tablets

Active ingredient: clobazam

Inactive ingredients: corn starch, lactose monohydrate, magnesium stearate, silicon dioxide, and talc.

Oral Suspension

Active ingredient: clobazam

Inactive ingredients: magnesium aluminum silicate, xanthan gum, citric acid monohydrate, disodium hydrogen phosphate dihydrate, simethicone emulsion, polysorbate 80, methylparaben, propylparaben, propylene glycol, sucralose, maltitol solution, berry flavor, purified water.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Marketed by:

Lundbeck, Deerfield, IL 60015, U.S.A.

Revised: 11/2013

Instructions for Use
ONFI® (ON-free)
(clobazam)
Oral Suspension

Read this Instructions for Use before using ONFI oral suspension and each time you get a refill. There may be new information. This leaflet does not take the place of talking with your healthcare provider about your medical condition or treatment.

Prepare ONFI Oral Suspension Dose

You will need the following supplies:

See Figure A

- ONFI oral suspension bottle
- Bottle adapter
- Oral dosing syringe (2 dosing syringes are included in the ONFI oral suspension box).
- Use only 1 syringe to take your dose of ONFI oral suspension. If you lose or damage the syringe, or cannot read the markings, use the other syringe.



Figure A

Step 1.

Remove the ONFI oral suspension bottle, bottle adapter, and 1 syringe from the box.

Step 2.

Shake the bottle well before each use.

See Figure B



Figure B

Step 3.

Uncap the bottle and firmly insert the bottle adapter into the bottle until the adapter top is even with the bottle top.

See Figure C

Once the bottle adapter is in place, it should not be removed.




Figure C

Step 4.

Check your dose in milliliters (mL) as prescribed by your healthcare provider. Find this number on the syringe. Do not take more than the prescribed total dose in 1 day.

See Figure D




Figure D

Step 5.

Push the plunger all the way down and then insert the syringe into the upright bottle through the opening in the bottle adapter.

See Figure E




Figure E

Step 6.

With the syringe in place, turn the bottle upside down. Pull the plunger to the number of mLs needed (the amount of liquid medicine in Step 4).

See Figure F




Figure F

Measure the mLs of medicine using the black ring on the white plunger.

See Figure G




Figure G

Step 7.

Remove the syringe from the bottle adapter. Slowly squirt ONFI oral suspension directly into the corner of your mouth or your child's mouth until all of the liquid medicine in the syringe is given.

See Figure H



Figure H

Step 8.

Cap the bottle tightly with the adapter in place. If the cap does not fit securely, check to see if the adapter is fully inserted.

See Figure I

Store the bottle upright at 68°F to 77°F (20°C to 25°C).

- Use ONFI oral suspension within 90 days of first opening bottle.
- After 90 days safely throw away any ONFI oral suspension that has not been used.



Figure I

Step 9.

Wash the oral syringe after each use.

- To clean the oral syringe, take apart by removing the plunger completely. Pull plunger straight out of the barrel.
- The barrel and plunger can be washed with soap and water, rinsed, and allowed to dry.
- Do not wash the oral syringe in the dishwasher.

Marketed by: Lundbeck, Deerfield, IL 60015, U.S.A.



ONFI is a registered trademark of Lundbeck

Revised: 11/2013

CLB-L-00005